

CERN-TH/96-363
 ICCR-Report-379-97-2
 hep-ph/9612493

Predictions of m_b/m_τ and m_t in an Asymptotically Non-Free Theory

MASAKO BANDO

*Aichi University, Miyoshi, Aichi 470-02, Japan**

TETSUYA ONOGI

Department of Physics, Hiroshima University, Hiroshima 606-01, Japan[†]

JOE SATO

Institute for Cosmic Ray Research

The University of Tokyo, Midori-Cho, Tanashi, Tokyo 188, Japan[‡]

TATSU TAKEUCHI

CERN, TH Division, CH-1211 Genève 23, Switzerland[§]

Abstract

We discuss an extension of the Minimal Supersymmetric Standard Model (MSSM) with 5 generations of matter superfields. The extra generations are assumed to form a generation-mirror generation pair (the 4th and anti-4th generations) enabling the extra fermions to have $SU(2)_L \times U(1)_Y$ invariant masses. Due to the contribution of the extra generations, all three running gauge couplings of $SU(3)_C \times SU(2)_L \times U(1)_Y$ become asymptotically non-free while preserving gauge coupling unification at the GUT scale. We show that due to the asymptotically non-free character of the gauge couplings: (1) the top and bottom Yukawa couplings are strongly focused onto infrared fixed points as they are evolved down in scale making their values at $\mu = M_Z$ insensitive to their initial values at $\mu = M_{\text{GUT}}$; (2) the model predicts $R_{b\tau}(M_Z) \equiv Y_b/Y_\tau|_{\mu=M_Z} \approx 1.8$, which is consistent with the experimental value provided we take the ratio of Yukawa couplings at the GUT scale to be $R_{b\tau}(M_{\text{GUT}}) = Y_b/Y_\tau|_{\mu=M_{\text{GUT}}} = 1/3$; (3) the t mass prediction comes out to be $m_t \approx 180 \text{ GeV}$ which is also consistent with experiment.

*E-mail: bando@aichi-u.ac.jp

[†]E-mail: onogi@theo.phys.sci.hiroshima-u.ac.jp

[‡]E-Mail: joe@icrhp3.icrr.u-tokyo.ac.jp

[§]E-mail: takeuchi@vxcern.cern.ch

1 Introduction

The popularity of the Minimal Supersymmetric Standard Model (MSSM) in recent years is mainly due to its success in attaining gauge coupling unification: given the particle content of the MSSM, the three coupling constants of the $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge groups converge to a common value at a common scale (the GUT scale) when evolved up to higher energies using the renormalization group equations (RGE) [1]. This unification of the gauge coupling constants is crucial if one wishes to construct a Grand Unified Theory (GUT) which unifies the three gauge groups of the Standard Model (SM) into a larger simple group at a single scale. However, it should be noted that the particle content which achieves such unification is not unique [2]. In particular, as pointed out in Ref. [3], one always has the freedom to add complete generations of matter superfields to the MSSM without destroying the unification condition.¹

Another attractive feature of the MSSM is the possibility of unifying the b and τ Yukawa couplings: if one assumes

$$R_{b\tau}(M_{\text{GUT}}) = Y_b(M_{\text{GUT}})/Y_\tau(M_{\text{GUT}}) = 1 \quad (1.1)$$

at the GUT scale², then one finds that the MSSM can reproduce the experimental value of $R_{b\tau}(M_Z) = Y_b(M_Z)/Y_\tau(M_Z) \approx 1.8$ if the Yukawa couplings of the top and the bottom were such that $Y_t(M_{\text{GUT}}) \gtrsim 2 \gg Y_b(M_{\text{GUT}})$ ($1 \lesssim \tan \beta \lesssim 3$), or $Y_t(M_{\text{GUT}}) \approx Y_b(M_{\text{GUT}}) \approx 1$ ($40 \lesssim \tan \beta \lesssim 60$).³ [5]

The reason why the experimental value of $R_{b\tau}$ can only be reproduced for either small or large $\tan \beta$ is easy to understand⁴: QCD interactions will enhance $Y_b(\mu)$ over $Y_\tau(\mu)$ as they are evolved down from M_{GUT} to M_Z so that $R_{b\tau}(M_Z)$ will end up well above the experimental value if only running due to gauge interactions were taken into account. This QCD effect must be countered by strong Yukawa interactions which will slow down the running. A smaller value of $R_{b\tau}$ consistent with experiment can be obtained when Y_t is large enough to counter the QCD enhancement alone, or when both Y_t and Y_b are large so that the two of them combined can have the desired effect. In the intermediate $\tan \beta$ region ($3 \lesssim \tan \beta \lesssim 40$) Y_t is not large enough to sufficiently suppress the increase of $R_{b\tau}$ by itself while Y_b is not large enough to compensate for it.

¹Of course, if one adds too many generations, the gauge couplings will reach the Landau pole before reaching the GUT scale. See Ref. [4].

²Whether the condition $Y_b(M_{\text{GUT}}) = Y_\tau(M_{\text{GUT}})$ is realized or not in GUT's depends on the representation of the Higgs field which gives mass to the fermions. For $SU(5)$, $SO(10)$, and E_6 unifications, the Higgs must be in the **5**, **10**, and **27** representations, respectively.

³Note that since $m_t/m_b = (Y_t/Y_b) \tan \beta$, the region $Y_t \gg Y_b$ corresponds to small $\tan \beta$ while $Y_t \approx Y_b$ corresponds to large $\tan \beta$. The lower and upper limits of $1 \lesssim \tan \beta$ and $\tan \beta \lesssim 60$ are required to keep Y_t and Y_b in the perturbative region throughout evolution between M_Z and M_{GUT} .

⁴We assume the reader has some familiarity with the RGE's for the Yukawa couplings

Of these two solutions, the small $\tan\beta$ case is often considered particularly attractive since the large size of $Y_t(M_{\text{GUT}})$ will drive $Y_t(\mu)$ rapidly towards an infrared *quasi*-fixed point [6] as it is evolved down in scale. As a result, the value of $Y_t(M_Z)$ is highly insensitive to its initial value $Y_t(M_{\text{GUT}})$ at the GUT scale. On the other hand, the large $\tan\beta$ case opens the possibility of unifying the top Yukawa coupling with the other two:

$$Y_t(M_{\text{GUT}}) = Y_b(M_{\text{GUT}}) = Y_\tau(M_{\text{GUT}}), \quad (1.2)$$

as required in $SO(10)$ unification with a **10**-Higgs. However, the insensitivity to the initial condition at M_{GUT} is lost.⁵

In this paper, we wish to outline how these conclusions will be modified when the MSSM is extended with an addition of a generation-mirror generation pair of extra matter superfields. (the 4th and $\bar{4}$ th generations)⁶. Each generation is assumed to consist of the usual 15 chiral fermion fields plus their superpartners. We will ignore the right-handed neutrino necessary to form the **16** representation of $SO(10)$ since we will always assume it to have a superheavy Majorana mass and make it decouple from the RGE's.⁷ Due to the mirror quantum number assignments between the 4th and $\bar{4}$ th generation fermions, they can develop $SU(2)_L \times U(1)_Y$ invariant masses enabling the left-handed neutrino to have a heavy Dirac mass thus circumventing the LEP limit for the number of massless neutrinos. Also, radiative corrections to LEP observables from the extra fermions can be made to decouple by making this gauge invariant mass large.[11]

One immediate consequence of the presence of the 2 extra generations is that all three gauge couplings of $SU(3)_C \times SU(2)_L \times U(1)_Y$ will be asymptotically non-free: they will become larger as they are evolved up to coincide at the unification scale [12]. This property is actually unique to the 5 generation model. In models with 4 generations⁸ or less, the QCD coupling will stay asymptotically free, and in models with 6 generations or more the couplings will diverge before unification.

As shown in the appendix, the unification of gauge couplings is controlled solely by the *differences* of the beta function coefficients in the one-loop approximation. Since the differences of the coefficients are independent of the number of full generations, the gauge coupling unification in our 5 generation model works well just as in the MSSM.

However, an important difference between asymptotically free theories and asymptotically non-free theories is that $\alpha = 0$ is an ultraviolet (UV) fixed point in the

⁵Another problem with the large $\tan\beta$ solution is that fine tuning of the Higgs potential is necessary to achieve radiative electroweak symmetry breaking. In the small $\tan\beta$ case, radiative electroweak symmetry breaking is naturally achieved due to the initial condition $Y_t(M_{\text{GUT}}) \gg Y_b(M_{\text{GUT}})$. However, fine tuning is necessary in this case also to obtain the correct value of $\tan\beta$. See, for instance, Ref. [7].

⁶ Such pairs are well known to exist in many GUT scenarios. See Ref. [8].

⁷We do not consider an intermediate scale for the right-handed neutrino mass for the sake of simplicity. See Refs. [9, 10] for analyses of the MSSM case with an intermediate scale.

⁸4 generation models have been discussed in Refs. [13].




Figure 1: The difference between asymptotically free and non-free theories.

former but an infrared (IR) fixed point in the latter. This means that for asymptotically free (non-free) theories, the RG flow will be such that a large region of α values in the IR (UV) will flow into a small region close to $\alpha = 0$ in the UV (IR), and the difference in the relative size of these regions will be more pronounced for larger separations in scale. Therefore, in order to get the desired value of $\alpha(M_Z)$ in asymptotically free theories, the value of $\alpha(M_{\text{GUT}})$ must be tuned to extreme accuracy while for asymptotically non-free theories, no fine tuning is necessary. This is shown schematically in Fig. 1.

This absence of the necessity to fine tune parameters at the UV cutoff is an extremely attractive feature of asymptotically non-free theories. It means that the high energy theory effective above the UV cutoff can give the correct predictions at low energies as long as it predicts the values of the running couplings at the cutoff to be within an only mildly restricted range. This point has been emphasized previously by many authors [14] (though not necessarily from a modern point of view). In particular, Moroi, Murayama and Yanagida [4] have studied the same 5 generation model as we are considering here and have shown that the values of the running couplings at M_{GUT} need not even be unified to predict the correct value of $\sin^2 \theta_w$.

In this paper, we extend the analysis of Moroi et al. and study how the existence of the extra generations will affect the running of the Yukawa couplings of the 3rd generation fermions. A similar problem for the non-supersymmetric case has been considered in [15]. As in the MSSM case, we will impose a unification condition on the Yukawa couplings at M_{GUT} and determine the parameter range in which our model can predict the correct top, bottom and τ -lepton masses.

The attentive reader at this point may think that such a program is doomed to failure from the beginning. Since the QCD coupling is asymptotically non-free, the QCD enhancement of $R_{b\tau}$ from M_{GUT} to M_Z will be even larger than the MSSM case making it impossible to bring $R_{b\tau}(M_Z)$ down to ~ 1.8 even with large Yukawa couplings. However, we would like to quickly point out that the unification condition need not be that of Eqs. (1.1) or (1.2). In fact, an $SO(10)$ -GUT with an **126**-Higgs predicts [16]

$$Y_t(M_{\text{GUT}}) = Y_b(M_{\text{GUT}}) = \frac{1}{3}Y_\tau(M_{\text{GUT}}), \quad (1.3)$$

so that $R_{b\tau}(M_{\text{GUT}}) = 1/3$. This is the unification condition which we will adopt.⁹ In this case, the extra enhancement from QCD is actually welcome since $R_{b\tau}$ must be enhanced by a factor of $5 \sim 6$ to reproduce the experimental value of $R_{b\tau}(M_Z)$.

This paper is organized as follows: In section 2, we describe our model and specify the way we calculate the RG evolution of the gauge and Yukawa couplings. In section 3, we show how the gauge couplings can be unified in our model. Section 4 discusses Yukawa coupling unification and the predictions for $R_{b\tau}(M_Z)$ and m_t . Section 5 concludes.

2 The $4 + \bar{1}$ Generation Model:

In extending the MSSM by introducing extra matter superfields, we must keep two things in mind: (1) the matter superfields must be introduced in such a way that gauge coupling unification (and anomaly cancellation) of the MSSM is preserved, and (2) the fermion content must be compatible with the constraints placed by LEP measurements, namely three massless neutrino species and the so-called Peskin–Takeuchi constraint [17].

The simplest way to satisfy these requirements is to introduce 2 extra generations which form a generation–mirror generation pair. We will call them the 4th and anti-4th generations. The fermion content of these extra generations will be ‘vector-like’ so that all of them, including the extra neutrinos, can develop $SU(2)_L \times U(1)_Y$ invariant Dirac masses. These masses will also suppress the size of radiative corrections to LEP observables from the extra fermions enabling them to circumvent the Peskin–Takeuchi constraint[11].

It should be noted that we can only introduce one such generation–mirror generation pair. If we introduce two pairs or more, all three gauge couplings will reach their Landau poles and diverge well before the would-have-been unification scale[4].

We denote the extra fermion families (U, D, N, E) and $(\bar{U}, \bar{D}, \bar{N}, \bar{E})$, respectively, and give them a common $SU(2)_L \times U(1)_Y$ invariant mass of M_{EVF} . Their superpartners, and all the other supersymmetric particles in the theory will be given

⁹An **126**-Higgs is necessary to give a direct Majorana mass term to the right-handed neutrino.

a common SUSY breaking mass of M_{SUSY} . For the sake of simplicity, we take $M_{\text{EVF}} = M_{\text{SUSY}} = 1 \text{ TeV}$.

In addition to the $SU(2)_L \times U(1)_Y$ invariant masses, we also couple the 4th and $\bar{4}$ th generation fermions to the two Higgs doublets in the same way as the other generations. Here we take the case where

$$\begin{aligned} Y_U &= Y_t, & Y_{\bar{U}} &= 0, \\ Y_D &= Y_b, & Y_{\bar{D}} &= 0, \\ Y_E &= Y_\tau, & Y_{\bar{E}} &= 0. \end{aligned} \quad (2.1)$$

and set all the 1st and 2nd generation Yukawa couplings to zero. Furthermore, we impose the unification condition

$$Y_t(M_{\text{GUT}}) = Y_b(M_{\text{GUT}}) = \frac{1}{3}Y_\tau(M_{\text{GUT}}) \equiv Y_{\text{GUT}}, \quad (2.2)$$

as mentioned in the introduction.

In view of the relatively large coupling strengths near the unification scale due to the asymptotically non-freeness, we use the fully coupled 2-loop renormalization group equations (RGE's) from Ref. [18] to evolve the gauge and Yukawa couplings. We ignore small differences in the masses of the 4th and $\bar{4}$ th generation particles or that of the supersymmetric particles which may be induced by the Yukawa couplings and simply set all their masses at $M_{\text{EVF}} = M_{\text{SUSY}} = 1 \text{ TeV}$. We also ignore threshold corrections. Therefore, between M_{GUT} and $M_{\text{EVF}} = M_{\text{SUSY}}$, we evolve the couplings with the RGE's for the Supersymmetric SM with 5 super-generations and 2 super-Higgs doublets, while between $M_{\text{EVF}} = M_{\text{SUSY}}$ and M_Z , we use the RGE's for the SM with only 3 ordinary generations and 1 Higgs doublet. The gauge couplings are connected continuously at $M_{\text{EVF}} = M_{\text{SUSY}} = 1 \text{ TeV}$ while the up-type (down-type) Yukawa couplings are multiplied by $\sin \beta$ ($\cos \beta$) below $M_{\text{EVF}} = M_{\text{SUSY}}$ to take into account the decoupling of one of the Higgses.

The number of adjustable parameters in our model is four: the unification scale M_{GUT} , the unified gauge coupling α_{GUT} , the unified Yukawa coupling Y_{GUT} , and the mixing angle of the low lying Higgs fields $\tan \beta = v_2/v_1$, where $\sqrt{v_1^2 + v_2^2} = v = 246 \text{ GeV}$. We restrict α_{GUT} and Y_{GUT} to the region

$$\alpha_{\text{GUT}} < 1.0, \quad Y_{\text{GUT}} < 0.7. \quad (2.3)$$

(Note that $Y_\tau(M_{\text{GUT}}) = 3Y_{\text{GUT}}$. Note also that the natural expansion coefficient corresponding to the $\alpha_i(\mu)$'s is $Y^2/(4\pi)$ for the Yukawa's.) As we will see later, this will keep the gauge and Yukawa couplings within their perturbative regions throughout the evolution from M_{GUT} to M_Z .

Since we do not consider the evolution of the soft SUSY breaking parameters of the Higgs potential in this paper, $\tan \beta$ will remain a phenomenological parameter to be fixed by hand. We will use the τ -lepton mass to fix $\tan \beta$ from

$$m_\tau(M_{\text{SUSY}}) = \frac{v}{\sqrt{2}}Y_\tau(M_{\text{SUSY}})\cos \beta. \quad (2.4)$$




Figure 2: The allowed region in the plane $(\alpha_{\text{GUT}}, M_{\text{GUT}})$. The small-black, and large-gray circles indicate the ranges $0.4 \leq Y_{\text{GUT}} < 0.7$, and $0.1 < Y_{\text{GUT}} < 0.4$, respectively.

3 Gauge Coupling Unification:

The values of the $SU(3)_C \times SU(2)_L \times U(1)_Y$ coupling constants at $\mu = M_Z$ are given by [19] :

$$\alpha_1(M_Z) = 0.01689 \pm 0.00005, \quad (3.1)$$

$$\alpha_2(M_Z) = 0.03322 \pm 0.00025, \quad (3.2)$$

$$\alpha_3(M_Z) = 0.12 \pm 0.01. \quad (3.3)$$

Note that these are the $\overline{\text{MS}}$ coupling constants¹⁰ and that the $U(1)$ coupling is normalized to $\alpha_1 = (5/3)g'^2/4\pi$.

For fixed values of Y_{GUT} in the range given in Eq. (2.3), we searched for values of α_{GUT} and M_{GUT} which reproduced the experimental data given above. The results are shown in Fig. 2.

We see that the allowed range of α_{GUT} is narrow for smaller M_{GUT} but still exist down to $M_{\text{GUT}} \approx 10^{16.55}\text{GeV}$ and becomes wider as M_{GUT} is increased. This result is as expected from our discussion on asymptotically non-free theories: a wider range of α_{GUT} corresponds to a much smaller range of couplings at M_Z , and the allowed range will become wider as M_{GUT} is increased. However, if we increase

¹⁰This may be a confusing point since the ‘effective’ QED coupling constant $\alpha(M_Z)$ and the ‘effective’ weak angle $\sin^2 \theta_W$ that are usually quoted by LEP are not $\overline{\text{MS}}$ values.




Figure 3: Typical μ dependence of $\alpha_1(\mu)$, $\alpha_2(\mu)$ and $\alpha_3(\mu)$. The parameter values for this plot were $(\alpha_{\text{GUT}}, Y_{\text{GUT}}, M_{\text{GUT}}, \tan \beta) = (0.35, 0.3, 10^{16.6} \text{GeV}, 57.5)$.

M_{GUT} beyond $\sim 10^{17.1} \text{GeV}$, then α_{GUT} and/or Y_{GUT} will have to be taken beyond the limits specified in Eq. (2.3) and they will be too large for the perturbative treatment of the RGE's to be reliable.

As an example, we show the running of the three gauge couplings in Fig. (3) for typical values of α_{GUT} , M_{GUT} , and Y_{GUT} . We see a small deviation from linear dependence on $\log \mu$ near M_{GUT} where the couplings become large and the two-loop corrections start contributing to the running appreciably. However it is clear that two-loop contributions are still not very serious within the range of α_{GUT} which we have chosen here and we may regard our perturbative treatment to be sufficient.

4 Yukawa Coupling Unification:

Next, we fix the values of α_{GUT} and M_{GUT} in the range allowed by gauge coupling unification and calculate the evolution of the Yukawa couplings for different values of Y_{GUT} . Typical evolutions of the τ , b , and t Yukawa couplings are shown in Figs. 4, 5, and 6. As is evident from these figures, the asymptotic non-freeness of the gauge couplings has a strong focusing effect on the top and bottom Yukawa couplings as they evolve down in scale and as a result, the values of the two Yukawa's converge to IR fixed points by the time they reach the SUSY breaking scale $M_{\text{SUSY}} = 1 \text{TeV}$. In the case of the τ Yukawa coupling, the situation is rather different. Near the GUT scale it tends to focus itself due to its larger size at M_{GUT} (Recall $Y_\tau(M_{\text{GUT}}) =$




Figure 4: The running of the τ -lepton Yukawa coupling for typical values of $(\alpha_{\text{GUT}}, M_{\text{GUT}})$ shown in Fig. 2. The value of α_{GUT} is varied between 0.3 and 0.8 while $Y_\tau(M_{\text{GUT}}) = 3Y_{\text{GUT}}$ is varied between 0.6 and 2.1. The Yukawa coupling is multiplied by $\cos \beta$ below M_{SUSY} .




Figure 5: The running of the bottom Yukawa coupling for typical values of $(\alpha_{\text{GUT}}, M_{\text{GUT}})$ shown in Fig. 2. The value of α_{GUT} is varied between 0.3 and 0.8 while $Y_b(M_{\text{GUT}}) = Y_{\text{GUT}}$ is varied between 0.2 and 0.7. The Yukawa coupling is multiplied by $\cos \beta$ below M_{SUSY} .




Figure 6: The running of the top Yukawa coupling for typical values of $(\alpha_{\text{GUT}}, M_{\text{GUT}})$ shown in Fig. 2. The value of α_{GUT} is varied between 0.3 and 0.8 while $Y_t(M_{\text{GUT}}) = Y_{\text{GUT}}$ is varied between 0.2 and 0.7. The Yukawa coupling is multiplied by $\sin \beta$ below M_{SUSY} .

$3Y_{\text{GUT}}$). However, unlike the top and bottom Yukawa couplings, once Y_τ becomes small at lower scales, it runs slowly and does not quite converge to its IR fixed point ($y_\tau = 0$).

Due to this IR fixed point behavior, the values of $Y_t(M_Z)$, $Y_b(M_Z)$ have almost no dependence on Y_{GUT} . They do depend on the value of α_{GUT} but even then only very mildly.

4.1 Bottom to Tau Mass Ratio:

The b - τ mass ratio has been the most intensively studied quantity in both supersymmetric and non-supersymmetric GUT scenarios. Many authors have investigated the possibility of unifying the two couplings with various degrees of success. [2, 5, 9, 10, 20, 21, 22, 23, 24]

Currently, the experimentally determined $\overline{\text{MS}}$ running masses of the τ -lepton and the b quark at $\mu = M_Z$ are given by [19]

$$\begin{aligned} m_\tau(M_Z) &= 1.75 \pm 0.01 \text{ GeV}, \\ m_b(M_Z) &= 3.1 \pm 0.4 \text{ GeV}, \end{aligned} \tag{4.1}$$

from which we conclude

$$R_{b\tau}(M_Z) = 1.6 \sim 2.0 \tag{4.2}$$




Figure 7: The dependence of $R_{b\tau}(M_Z)$ on M_{GUT} and α_{GUT} . The small-black, and large-gray circles indicate the ranges $0.3 < \alpha_{\text{GUT}} \leq 0.6$, and $0.6 < \alpha_{\text{GUT}} < 1$, respectively. The dependence of $R_{b\tau}(M_Z)$ on Y_{GUT} is negligible.

The dependence of $R_{b\tau}(M_Z)$ on α_{GUT} and M_{GUT} in our model is shown in Fig. 7. Obviously, whether our model can reproduce the experimental value of $R_{b\tau}(M_Z)$ or not depends almost solely on the value of α_{GUT} . If $\alpha_{\text{GUT}} > 0.6$, then the QCD interactions near M_{GUT} will be so strong that Y_b will be enhanced too much relative to Y_τ . However, there is still a large set of $(\alpha_{\text{GUT}}, M_{\text{GUT}})$ values which keeps $R_{b\tau}(M_Z)$ below 2.

Of course, since $\tan \beta$ is large in our model there is potentially a very large radiative correction to m_b from the loop induced coupling of the b quark to v_2 . [2, 22, 25, 26] This correction can throw our prediction off the mark by a considerable amount.

However, such corrections can easily be compensated for. If the correction makes $R_{b\tau}(M_Z)$ smaller, we only need to make α_{GUT} larger. If it makes $R_{b\tau}(M_Z)$ larger, we only need to make α_{GUT} smaller, changing M_{EVF} and/or M_{SUSY} if necessary.

We therefore conclude that our model can accommodate the b - τ mass ratio quite easily without any fine tuning.

4.2 The Top Quark Mass:

Due to the IR fixed point behavior of Y_τ , Y_b , and Y_t we have seen above, for the range of $(\alpha, M_{\text{GUT}}, Y_{\text{GUT}})$ values that yield the correct value of $R_{b\tau}(M_Z)$ we find:

$$Y_t(M_Z) = 1.01 \sim 1.07. \quad (4.3)$$

Using the τ mass to fix $\tan\beta$, we find

$$m_t(M_Z) = 176 \sim 187 \text{ GeV}, \quad (4.4)$$

which is in perfect agreement with the experimental value [19] :

$$m_t(M_Z) = 180 \pm 10 \text{ GeV}. \quad (4.5)$$

This result is actually highly dependent on our choice Eq. (2.1) for the 4th and $\bar{4}$ th generation Yukawa couplings. Had we chosen a different condition such as

$$\begin{aligned} Y_U &= Y_{\bar{U}} = Y_t, \\ Y_D &= Y_{\bar{D}} = Y_b, \\ Y_E &= Y_{\bar{E}} = Y_\tau, \end{aligned} \quad (4.6)$$

then the IR fixed value for Y_t would have been

$$Y_t(M_Z) \sim 0.705 \quad (4.7)$$

leading to a prediction of

$$m_t(M_Z) \sim 154 \text{ GeV} \quad (4.8)$$

5 Summary and Conclusions :

We have presented an extension of the MSSM with a generation anti-generation pair of extra matter superfields. The $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge couplings are asymptotically non-free in this model but still converge to a common value before any of them diverge. Consequences of this asymptotically non-free theory are:

1. The unified coupling α_{GUT} and the unification scale M_{GUT} need not be fine tuned to reproduce the values of the $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge couplings at $\mu = M_Z$. Larger values of M_{GUT} allow for a larger range of α_{GUT} values. However, M_{GUT} must be taken below $\sim 10^{17.1} \text{ GeV}$ to keep α_{GUT} and Y_{GUT} within perturbative range.
2. The top and bottom Yukawa couplings are strongly focused onto IR fixed points. This makes the IR values of the two Yukawa couplings insensitive to their initial value Y_{GUT} at the GUT scale.

3. The model can reproduce the ratio of bottom and tau masses $R_{b\tau}(M_Z) = Y_b(M_Z)/Y_\tau(M_Z) = 1.6 \sim 2$, provided we assume the unification condition $R_{b\tau}(M_{\text{GUT}}) = 1/3$ (**126**-Higgs) instead of the usual $R_{b\tau}(M_{\text{GUT}}) = 1$ (**10**-Higgs).
4. The model can also give the correct top mass provided the Yukawa couplings of the 4th and $\bar{4}$ th generations are taken as in Eq. (2.1).

It would be most interesting if this insensitivity of the low energy predictions to the initial conditions at the GUT scale could be taken further to include the soft SUSY breaking parameters of the Higgs sector. In particular, if a large $\tan\beta$ could be generated without fine tuning, it could provide an answer to the question why the top is so much heavier than the bottom.

Above M_{GUT} , our model is a supersymmetric $SO(10)$ theory which includes four **16**-plets and an $\bar{\mathbf{16}}$ -plet, and a Higgs sector which consist of at least an **126** and an $\bar{\mathbf{126}}$ (in order for **126** to have a mass term). This makes the $SO(10)$ gauge coupling asymptotically non-free also.

It has recently been argued that such asymptotic non-freeness of the gauge couplings can be consistently interpreted as a sign of compositeness. [27] The basic reasoning is as follows: The general compositeness condition of gauge bosons is given by $Z(M_{\text{comp}}) = 0$ where $Z(\mu)$ is the wave-function renormalization constant and M_{comp} is the compositeness scale. If one enforces conventional normalization $Z(\mu) = 1$ at all scales μ , then superficially the running gauge coupling $\alpha(\mu)$ will diverge at $\mu = M_{\text{comp}}$ making it look like an asymptotically non-free theory. This is analogous to theories with dynamically generated Higgs bosons where compositeness manifests itself as the divergence of the Yukawa coupling at the compositeness scale in the effective Higgs-Yukawa theory [28].

In the SM, the large number of arbitrary parameters have lead most people to believe that going beyond the SM will somehow reduce the number of parameters and make theories more predictive. However, most extensions of the SM such as the MSSM or Technicolor actually *increases* the number of parameters by a huge amount. Reduction of the number of parameters is usually achieved by imposing ad hoc symmetries such as R -parity, universality of the scalar masses at the GUT scale, etc. What our analysis has shown is that for certain types of theories with IR fixed points, it may happen that most of the parameters simply *do not matter* or only needs to be specified to an order of magnitude to make precise low energy predictions. Clearly, the IR fixed point phenomena is an alternative to symmetries for making theories more predictive and deserves thorough and systematic investigation.

Acknowledgments

We would like to thank T. Kugo and T. Yanagida for their valuable comments and encouragement. Helpful discussions with W. A. Bardeen, M. Carena, K.-I. Izawa,

T. Takahashi, Y. Taniguchi, S. Tanimura, and C. E. M. Wagner are also gratefully acknowledged. The early stage of this work began when one of the authors (M.B.) was staying at Fermilab as a summer visitor in 1995 to collaborate with T.O. and T.T.. We would like to thank the members of the Fermilab Theory Group for providing us with this opportunity and for their warm hospitality during our stay. The main part of this work was completed during the Kashikōjima Workshop held in August 1996 at the Kashikōjima Center, Agō-chō, Shima-gun, Mié-ken, Japan, which was supported by the Grant-in Aid for Scientific Research (No. 07304029) from the Japanese Society for the Promotion of Science (JSPS). M.B. is supported in part by the Grant-in Aid for Scientific Research (No. 33901-06640416) also from the JSPS.

A Solution to the one-loop RG equations

In this appendix, we give a qualitative interpretation of our results in the one-loop approximation.

The one-loop renormalization group equations above the SUSY scale in our model are as follows:

$$\frac{d\alpha_i}{dt} = \frac{b_i}{2\pi}\alpha_i^2 \quad (i = 1, 2, 3) \quad (\text{A.1})$$

$$\frac{d\alpha_t}{dt} = \frac{\alpha_t}{2\pi} \left[-\left(\frac{13}{15}\alpha_1 + 3\alpha_2 + \frac{16}{3}\alpha_3 \right) + (9\alpha_t + \alpha_b) \right] \quad (\text{A.2})$$

$$\frac{d\alpha_b}{dt} = \frac{\alpha_b}{2\pi} \left[-\left(\frac{7}{15}\alpha_1 + 3\alpha_2 + \frac{16}{3}\alpha_3 \right) + (\alpha_t + 9\alpha_b + 2\alpha_\tau) \right] \quad (\text{A.3})$$

$$\frac{d\alpha_\tau}{dt} = \frac{\alpha_\tau}{2\pi} \left[-\left(\frac{9}{5}\alpha_1 + 3\alpha_2 \right) + (6\alpha_b + 5\alpha_\tau) \right]. \quad (\text{A.4})$$

Here, $t = \log \mu$, $\alpha_{t,b,\tau} \equiv \frac{Y_{t,b,\tau}^2}{4\pi}$, and the one-loop beta function coefficients are given by the following formula:

$$\begin{aligned} (b_1, b_2, b_3) &= (0, -6, -9) + (3 + 2n_{\text{vector}})(2, 2, 2) + 2\left(\frac{3}{10}, \frac{1}{2}, 0\right) \\ &\quad \text{(for the MSSM plus } n_{\text{vector}} \text{ full generation pairs)} \\ &= \left(\frac{53}{5}, 5, 1\right) \quad \text{(for } n_{\text{vector}} = 1\text{)}. \end{aligned}$$

Since the model is nothing but the standard model below the SUSY scale, as can be seen from Fig. 3, the experimental inputs are essentially equivalent to

$$\alpha_1^{-1}(t_{\text{SUSY}}) \approx 58, \quad \alpha_2^{-1}(t_{\text{SUSY}}) \approx 31, \quad \alpha_3^{-1}(t_{\text{SUSY}}) \approx 12.$$

The solution to Eq.(A.1) is

$$\frac{1}{\alpha_i(t)} = \frac{b_i}{2\pi}(t_{\text{GUT}} - t) + \frac{1}{\alpha_i(t_{\text{GUT}})} \quad (i = 1, 2, 3).$$

From the above equation, and using $\alpha_1^{-1}(t_{\text{SUSY}})$ and $\alpha_2^{-1}(t_{\text{SUSY}})$ as inputs, we can predict $\alpha_3^{-1}(t_{\text{SUSY}})$ as

$$\alpha_3^{-1}(t_{\text{SUSY}}) = \frac{b_3 - b_2}{b_1 - b_2} [\alpha_1^{-1}(t_{\text{SUSY}}) - \alpha_2^{-1}(t_{\text{SUSY}})] + \alpha_2^{-1}(t_{\text{SUSY}}) \approx 12.$$

Note that only the differences of beta function coefficients appear in this expression. Therefore, at the one-loop level, the prediction of $\alpha_3^{-1}(t_{\text{SUSY}})$ would be exactly the same for any n_{vector} . However, the two-loop correction which is $O(\max(\alpha^2(t)))$ would be different depending on whether the theory is asymptotically free or non-free. In the MSSM, the expected correction would be as large as $O(\alpha^2(t_{\text{SUSY}})) \sim 1\%$, whereas in the $n_{\text{vector}} = 1$ case, the correction would be as large as $O(\alpha^2(t_{\text{GUT}})) \sim 10\%$. Thus

if the one-loop prediction differs from the experiment by more than a few percent, one has to consider rather large threshold correction at the GUT scale to explain the discrepancy in the MSSM whereas in the latter model there is still room for two-loop corrections to account for it.

Next, let us solve Eqs. (A.2)~(A.4) to obtain the low energy behavior of the Yukawa couplings. The first term in each equation is the contribution from the gauge interactions and the second term in each equation is that from the Yukawa interactions. The gauge interactions try to enhance the Yukawa couplings as the scale is lowered whereas the Yukawa interactions tend to reduce it. Therefore, one can naively expect that the Yukawa couplings fall into infrared fixed points where the gauge contribution and the Yukawa contribution balance. Whether this is true or not depends on initial values and the details of the beta function coefficients.

In order to see this more explicitly, let us make the following three assumptions and reduce Eqs. (A.2)~(A.4) into a much simpler form:

1. Since at low energy, $\alpha_1, \alpha_2, \ll \alpha_3$, we can neglect α_1, α_2 .
2. At low energy, the contribution from α_τ is not so dominant compared to those from $\alpha_{t,b}$. This is partly because the coefficient of α_τ in Eq. (A.3) is not so large and partly because in Eq. (A.4) there is no contribution from α_3 which can prevent α_{yukawa} getting small thus α_τ gets small at lower energy much faster than $\alpha_{t,b}$.
3. Assuming 1 and 2, the difference between α_t and α_b is almost negligible. This is because we impose $\alpha_t(t_{\text{GUT}}) = \alpha_b(t_{\text{GUT}})$ as the GUT scale initial condition, and because the approximate RG equation is symmetric under the interchange $\alpha_t \leftrightarrow \alpha_b$.

In the following, we will obtain the solutions to the resulting approximate equations. Of course, the behavior of those solutions will be correct only qualitatively since the corrections from the neglected terms are not completely negligible. (In principle, it is possible to check the validity of this approximation by solving the full equations.) However, since we are only interested in the qualitative features, we will not discuss the corrections from the neglected terms here.

With these assumptions and by setting $\alpha_t = \alpha_b \equiv \alpha_Q$, Eqs. (A.2), (A.3) become

$$\frac{d\alpha_Q}{dt} = \frac{\alpha_Q}{2\pi} \left(-\frac{16}{3}\alpha_3 + 10\alpha_Q \right). \quad (\text{A.5})$$

Let us define the ratio $z_Q = \frac{\alpha_Q}{\alpha_3}$. Eq. (A.5) is then

$$\frac{dz_Q}{dt} = \frac{z_Q}{2\pi} \alpha_3 \left(-\frac{19}{3} + 10z_Q \right). \quad (\text{A.6})$$

It is easy to see that the solution to Eq. (A.6) is

$$\frac{z_Q(t) - \frac{19}{30}}{z_Q(t)} = \frac{z_Q(t_{\text{GUT}}) - \frac{19}{30}}{z_Q(t_{\text{GUT}})} \exp \left[-\frac{19}{3} \log \left(\frac{\alpha_3(t_{\text{GUT}})}{\alpha_3(t)} \right) \right] \quad (\text{A.7})$$

The exponent $\frac{19}{3} \log \left(\frac{\alpha_3(t_{\text{GUT}})}{\alpha_3(t)} \right)$ at $t = t_{\text{SUSY}}$ ranges roughly from 8 to 12, thus at t_{SUSY} the quark Yukawa coupling gets very close to the infrared fixed point.

$$\alpha_Q(t) \xrightarrow{t \rightarrow t_{\text{SUSY}}} \frac{19}{30} \alpha_3(t_{\text{SUSY}}) \quad (\text{A.8})$$

This behavior of the quark Yukawa coupling is consistent with what we see from the numerical solution to the two-loop RG equation.

On the other hand, the one-loop RG equation for the bottom-tau ratio $R_{b\tau}$ can be obtained from Eqs. (A.3), (A.4). Using the assumptions, the equation at low energy can be simplified to

$$\frac{dR_{b\tau}}{dt} = \frac{R_{b\tau}}{2\pi} \left(-\frac{8}{3} \alpha_3 + 2\alpha_Q \right) \approx \frac{R_{b\tau}}{2\pi} \left(-\frac{7}{5} \alpha_3 \right). \quad (\text{A.9})$$

It is easy to see that the solution to Eq. (A.9) is

$$R_{b\tau}(t) = R_{b\tau}(t_{\text{GUT}}) \exp \left[\frac{7}{5} \log \left(\frac{\alpha_3(t_{\text{GUT}})}{\alpha_3(t)} \right) \right] \quad (\text{A.10})$$

The factor $\exp \left[\frac{7}{5} \log \left(\frac{\alpha_3(t_{\text{GUT}})}{\alpha_3(t)} \right) \right]$ at $t = t_{\text{SUSY}}$ is about 4.1 to 7.5 for $\alpha_{\text{GUT}} = 0.3 \sim 0.55$. This gives roughly the right enhancement for $R_{b\tau}$.

On the other hand, the RG equations in the MSSM are given by:

$$\frac{d\alpha_i}{dt} = \frac{b_i}{2\pi} \alpha_i^2 \quad (i = 1, 2, 3) \quad (\text{A.11})$$

$$\frac{d\alpha_t}{dt} = \frac{\alpha_t}{2\pi} \left[-\left(\frac{13}{15} \alpha_1 + 3\alpha_2 + \frac{16}{3} \alpha_3 \right) + (6\alpha_t + \alpha_b) \right] \quad (\text{A.12})$$

$$\frac{d\alpha_b}{dt} = \frac{\alpha_b}{2\pi} \left[-\left(\frac{7}{15} \alpha_1 + 3\alpha_2 + \frac{16}{3} \alpha_3 \right) + (\alpha_t + 6\alpha_b + \alpha_\tau) \right] \quad (\text{A.13})$$

$$\frac{d\alpha_\tau}{dt} = \frac{\alpha_\tau}{2\pi} \left[-\left(\frac{9}{5} \alpha_1 + 3\alpha_2 \right) + (3\alpha_b + 4\alpha_\tau) \right]. \quad (\text{A.14})$$

The one-loop beta function coefficients are given by the following formula:

$$(b_1, b_2, b_3) = \left(\frac{33}{5}, 1, -3 \right).$$

The equation for z_Q is

$$\frac{dz_Q}{dt} = \frac{z_Q}{2\pi} \alpha_3 \left(-\frac{19}{3} + 7z_Q \right). \quad (\text{A.15})$$

It is easy to see that the solution to Eq. (A.15) is

$$\frac{z_Q(t) - \frac{19}{21}}{z_Q(t)} = \frac{z_Q(t_{\text{GUT}}) - \frac{19}{21}}{z_Q(t_{\text{GUT}})} \exp \left[-\frac{19}{9} \log \left(\frac{\alpha_3(t)}{\alpha_3(t_{\text{GUT}})} \right) \right] \quad (\text{A.16})$$

The exponent $\frac{19}{9} \log \left(\frac{\alpha_3(t)}{\alpha_3(t_{\text{GUT}})} \right)$ at $t = t_{\text{SUSY}}$ is roughly 2, thus at t_{SUSY} the focusing effect of quark Yukawa coupling is not as strong as in our 5 generation model.

References

- [1] U. Amaldi, W. de Boer and H. Fürstenau, Phys. Lett. **B260** (1991) 447,
P. Langacker and N. Polonsky, Phys. Rev. **D52** (1995) 3081 [hep-ph/9503214].
- [2] A. Giveon, L. J. Hall, and U. Sarid,
Phys. Lett. **B271** (1991) 138, [KEK Scanned Preprint 9109088]
L. J. Hall, R. Rattazzi, and U. Sarid,
Phys. Rev. **D50** (1994) 7048 [hep-ph/9306309],
R. Rattazzi and U. Sarid,
Phys. Rev. **D53** (1996) 1553 [hep-ph/9505428],
U. Sarid, UND-HEP-96-US02 [hep-ph/9610341].
- [3] L. Ibáñez and G. Ross, Phys. Lett. **B105** (1981) 439,
W. J. Marciano and G. Senjanovic, Phys. Rev. **D25** (1981) 3092.
- [4] T. Moroi, H. Murayama and T. Yanagida,
Phys. Rev. **D48** (1993) R2995 [hep-ph/93062268].
- [5] P. Langacker and N. Polonsky,
Phys. Rev. **D49** (1994) 1454 [hep-ph/9306205];
Phys. Rev. **D50** (1994) 2199 [hep-ph/9403306],
N. Polonsky, Phys. Rev. **D54** (1996) 4537 [hep-ph/9602206].
- [6] B. Pendleton and G. G. Ross, Phys. Lett. **B98** (1981) 291,
C. T. Hill, Phys. Rev. **D24** (1981) 691.
- [7] M. Bando, T. Kugo, N. Maekawa, and H. Nakano,
Mod. Phys. Lett. **A7** (1992) 3379 [KEK Scanned Preprint 9212293].
M. Bando, N. Maekawa, H. Nakano, and J. Sato,
Mod. Phys. Lett. **A8** (1993) 2729 [KEK Scanned Preprint 9405421].
- [8] H. Georgi, Nucl. Phys. **B156** (1979) 126.
- [9] F. Vissani and A. Y. Smirnov, Phys. Lett. **B341** (1994) 173 [hep-ph/9405399].
- [10] A. Brignole, H. Murayama and R. Rattazzi,
Phys. Lett. **B335** (1994) 345 [hep-ph/9406397].
- [11] L. Lavoura and J. P. Silva,
Phys. Rev. **D47** (1993) 2046, [KEK Scanned Preprint 9302379]
N. Maekawa, Prog. Theor. Phys. **93** (1995) 919 [hep-ph/9406375];
Phys. Rev. **D52** (1995) 1684 [KEK Scanned Preprint 9504209];
KUNS-1366 [hep-ph/9510414].
- [12] S. Theisen, N.D. Tracas and G. Zoupanos, Z. Phys. **C37** (1988) 597

- [13] J. Bagger, S. Dimopoulos, and E. Masso, Nucl. Phys. **B253** (1985) 397,
 J. E. Björkman and D. R. T. Jones, Nucl. Phys. **B259** (1985) 533,
 M. Cvetič and C. R. Preitschopf, Nucl. Phys. **B272** (1986) 490,
 J. F. Gunion, D. W. McKay, and H. Pois,
 Phys. Lett. **B334** (1994) 339 [hep-ph/9406249],
 Phys. Rev. **D53** (1996) 1616 [hep-ph/9507323],
 M. Carena, H. E. Haber, and C. E. M. Wagner,
 Nucl. Phys. **B472** (1996) 55 [hep-ph/9512446].
- [14] G. Parisi, Phys. Rev. **D11** (1975) 909,
 L. Maiani, G. Parisi and R. Petronzio, Nucl. Phys. **B136** (1978) 115,
 N. Cabibbo and C.R. Farrar, Phys. Lett. **B110** (1982) 107.
- [15] G. Grunberg,
 Phys. Rev. **D38** (1988) R1012;
 Phys. Lett. **B203** (1988) R413.
- [16] M. S. Chanowitz, J. Ellis and M. K. Gaillard,
 Nucl. Phys. **B128** (1977) 506.
- [17] M. E. Peskin and T. Takeuchi,
 Phys. Rev. Lett. **65** (1990) 964, Phys. Rev. **D46** (1992) 381,
 J. L. Hewett, T. Takeuchi, and S. Thomas,
 SLAC-PUB-7088, CERN-TH/96-56 [hep-ph/9603391].
- [18] V. Barger, M. S. Berger, and P. Ohmann,
 Phys. Rev. **D47** (1993) 1093 [hep-ph/9209232],
 S. P. Martin and M. T. Vaughn, Phys. Lett. **B318** (1993) 331 [hep-ph/9308212].
- [19] Particle Data Group, Phys. Rev. **D54** (1996) 1.
- [20] S. Kelley, J. L. Lopez and D. V. Nanopoulos,
 Phys. Lett. **B274** (1992) 387 [KEK Scanned Preprint 9111309].
- [21] V. Barger, M. S. Berger, P. Ohmann, and R. J. N. Phillips,
 Phys. Lett. **B314** (1993) 351 [hep-ph/9304295];
 Phys. Rev. **D51** (1995) 2438 [hep-ph/9407273].
- [22] W. A. Bardeen, M. Carena, S. Pokorski and C. E. M. Wagner,
 Phys. Lett. **B320** (1994) 110 [hep-ph/9309293].
 M. Carena, M. Olechowski, S. Pokorski, and C. E. M. Wagner,
 Nucl. Phys. **B419** (1994) 213 [hep-ph/9311222];
 Nucl. Phys. **B426** (1994) 269 [hep-ph/9402253].
- [23] M. Bando, K.-I. Izawa and T. Takahashi,
 Prog. Theor. Phys. **92** (1994) 143 [hep-ph/9403284];
 Prog. Theor. Phys. **92** (1994) 1137 [hep-ph/9408314].

- [24] M. Olechowski and S. Pokorski,
 Phys. Lett. **B344** (1995) 201 [hep-ph/9407404],
 F. M. Borzumati, M. Olechowski, and S. Pokorski,
 Phys. Lett. **B349** (1995) 311 [hep-ph/9412379],
 H. Murayama, M. Olechowski and S. Pokorski,
 Phys. Lett. **B371** (1996) 57 [hep-ph/9510327].
- [25] T. Banks, Nucl. Phys. **B303** (1988) 172.
- [26] R. Hempfling, Phys. Rev. **D49** (1994) 6168.
- [27] M. Bando, Y. Taniguchi and S. Tanimura,
 KU-AMP 96014, KUNS-1420, HE(TH) 96/15 [hep-th/9610244].
- [28] W.A. Bardeen, C.T. Hill and M. Lindner, Phys. Rev. **D41** (1990) 1647.